

# DECON-QUAT® 200C

Disinfectant • Non-Food Contact Sanitizer • Mildewstat • Fungicide • Virucide\*

## ACTIVE INGREDIENTS:

Didecyl Dimethyl Ammonium Chloride.....10.14%  
Alkyl (50% C14, 40% C12, 10% C16)

Dimethyl Benzyl Ammonium Chloride.....6.76%

OTHER INGREDIENTS:.....83.10%

TOTAL:.....100.00%

**KEEP OUT OF REACH OF CHILDREN  
DANGER PELIGRO**

See side panel for additional precautionary statements.

## FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

For chemical or environmental emergencies, call CARECHEM 24 at 866-928-0739.  
Case Label: DQ200C-00-SM-NS Rev 6/16 • A-16 • Non-Sterile

**Vetel Associates, Inc.**

15 Lee Blvd. Malvern, PA 19355-1234

Tel: 610-644-8335 • Fax: 610-644-8336

www.sterile.com • US and Foreign Patents, www.sterile.com/patents.

EPA Reg. No. 10324-141-68959 • EPA Est. No. 68959-PA-01

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product can be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

Use	Dilution	Contact Time
For Hospital or Medical Environment claims	0.5 oz. /gal. water	10 minutes
For General or Broad Spectrum claims	0.5 oz. /gal. water	10 minutes
For Public Health Virucidal* claims	0.5 oz. /gal. water	10 minutes
For Non-Food Contact Sanitizing claims	0.5 oz. /gal. water	3 minutes
For Fungicidal claims	0.5 oz. /gal. water	10 minutes
For Mold and Mildew claims	0.5 oz. /gal. water	10 minutes

**FOR USE AS A ONE-STEP, GENERAL, HOSPITAL, MEDICAL DISINFECTANT, FUNGICIDE, VIRUCIDE\*, CLEANER:**

1. Pre-clean heavily soiled areas.
2. Prepare a solution according to SIMPLEX® System Container instructions to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray.
3. Treated surfaces must remain wet for 10 minutes.
4. Wipe dry with a clean cloth or allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished.
5. Prepare a fresh solution daily or when visibly dirty.

## BLOODBORNE PATHOGEN INSTRUCTIONS

**\*KILLS HIV, HBV AND HCV ON PRE-CLEANED HARD, NON-POROUS SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**  
**Personal Protection:** Wear protective latex gloves, gowns, masks and eye protection.  
**Cleaning Procedure:** Blood and other body fluids containing HIV-1, HBV & HCV must be thoroughly cleaned from hard, non-porous surfaces and objects before application of this product. **Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **Contact Time:** Allow hard, non-porous surfaces to remain wet for 10 minutes for all organisms.

**TO CONTROL MOLD/MILDEW:** Pre-clean heavily soiled areas. Prepare a solution according to SIMPLEX® System Container instructions. Apply use solution to hard, non-porous surfaces for 10 minutes which will effectively inhibit the growth of mold and mildew and their odors. Repeat treatment every seven days, or more often if new growth appears.

## NON-FOOD CONTACT SANITIZING DIRECTIONS

**NON-FOOD CONTACT SANITIZING:** Pre-clean heavily soiled areas. Prepare a solution according to SIMPLEX® System Container instructions. Apply solution to hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device, hand pump trigger spray device, coarse trigger spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Treated surfaces must remain wet for 3 minutes. Prepare a fresh solution daily or when visibly dirty.

**GENERAL DEODORIZATION:** To deodorize, apply 0.5 to 1 oz. of this product per gal. of water to hard, non-porous surfaces. Allow to air dry.

## AUTOMATED DILUTION SYSTEM (SIMPLEX® SYSTEM CONTAINER) INSTRUCTIONS:

### SIMPLEX® SYSTEM CONTAINER

Trigger Spray Bottle lid label:

1. To prepare use solution, open cap.
2. Peel off inner seal by grasping tab at far edge and pulling off.
3. Firmly push small, inner container all the way down.
4. Replace cap and tighten.
5. Slowly swirl for 15 seconds.
6. Move spray nozzle to open position.
7. Follow directions for use on label.

### SIMPLEX® SYSTEM CONTAINER

Gallon Size Bottle lid label:

1. To prepare use solution, open cap.
2. Peel off inner seal by grasping tab at far edge and pulling off.
3. Firmly push small, inner container all the way down.
4. Replace cap and tighten.
5. Slowly swirl for 15 seconds.
6. Open small side spout and peel off inner seal, as above.
7. Pour solution from small side spout onto surfaces to be treated or alternate containers.
8. Follow directions for use on label.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container 1/4 full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.



Made in the USA  
Fabriqué aux États-Unis  
Hecho en los EE. UU.

