



Medical

Pall Lipipor™ TNA Filter Set



Air-eliminating filter set for lipid-containing nutritional IV administration

Features

- ▶ Pall Lipipor 1.2 µm nylon membrane
- ▶ Smooth housing design
- ▶ Sterile, non-pyrogenic fluid pathway
- ▶ Phthalate-free tubing extension

Benefits

- ▶ Removes particles and microorganisms
- ▶ Retains oversized lipid droplets
- ▶ Eliminates air (independent of filter position)
- ▶ Easy to prime and use
- ▶ Increases patient comfort

Specifications

Filter Media

1.2 µm Pall Lipipor™ Nylon membrane

Priming Volume

1.5 mL

Sterility

Sterile and non-pyrogenic fluid pathway

Tubing Extension

Phthalate-free

Maximum Recommended Flow Rate

300 mL/hour


Maximum Working Pressure

Approximately 22 psi (1.5 bar)

Product Features

The Pall Lipipor TNA filter set is an air-elimination filter for use with any nutritional IV administration containing lipids and lipid emulsions. It is indicated for the removal of particulate debris, microbial contaminants, and entrained air that may be found in solutions intended for IV administration. It provides patient protection against particulate contamination, oversized lipid droplets¹, microorganisms^{2,3}, and air⁴.


Contamination of Parenteral Nutrition Preparations Can Have Serious Consequences



- Particulate contamination arises from infusion systems and components due to manipulations⁵, and interactions between components⁶. Particles can be deposited in the microvasculature of the lungs and other organs,

resulting in serious clinical consequences^{7,8}. Gross precipitation in admixtures, which may not be visible when lipid is present, has proven fatal in cases⁶.

- Oversized lipid droplets arise in admixtures due to instability. Large numbers may lodge in the lung microvasculature and produce an embolic syndrome¹.



- Microbial contamination can inadvertently arise in infusion systems due to manipulations. Parenteral nutrition is an acknowledged risk for fungemia with *Candida spp.* being among the most common organisms involved⁸. *Malassezia furfur* is also

emerging as an increasingly important pathogen in neonates³, having demonstrated the ability to grow in lipid-containing preparations^{3,9}.

Ordering Information

Description	Reorder Code		Packaging	
	USA	Europe	USA	Europe
Pall Lipipor TNA Filter for Parenteral Nutrition	TNA1	TNA1E	40/case	50/case

References

1. Driscoll, D.F., Bacon, M.N., & Bistran, B.R. Effects of in-line filtration on lipid particle size distribution in total nutrient admixtures. *JPEN J Parenter Enteral Nutr.* 1996; 20:296-301.
2. Barnett, M.I. Filtration of lipid-containing total parenteral nutrition (TPN) admixture. *Clin Nutr.* 1995; 14:49.
3. Robinson, R. & Ball, P. Presented at the New Zealand Hospital Pharmacists Association Meeting. October, 1996, Auckland, NZ.
4. Coppa, G.F., Gouge, T.H., & Hofstetter, S.R. Air embolism: a lethal but preventable complication of subclavian vein catheterization. *JPEN J Parenter Enteral Nutr.* 1981; 5:166-8.
5. Foroni, L.A., Rochat, M.H., Trouiller, P., & Calop, J.Y. Particle contamination in a ternary nutritional admixture. *J Parenter Sci Technol.* 1993; 47:311-4.
6. United States Food & Drug Administration Safety Alert: Hazards of precipitation associated with parenteral nutrition. April 18, 1994.
7. Walpot, H., Franke, R.P., Burchard, W.G., Agternkamp, C., Muller, F.G., & Mittermayer, C., et al. Particulate contamination of infusion solutions and drug additives within the scope of long-term intensive therapy. *Anaesthesist.* 1989; 38:544-8.
8. Lehr, H.A., Brunner, J., Rangoonwala, R., & Kirkpatrick, C.J. Particulate matter contamination of intravenous antibiotics aggravates loss of functional capillary density in postischemic striated muscle. *Am J Respir Crit Care Med.* 2002; 165:514-20.
9. Vazquez, J.A., Sanchez, V., Dmuchowski, C., Dembry, L.M., Sobel, J.D., & Zervos, M.J. Nosocomial acquisition of *Candida albicans*: an epidemiologic study. *J Infect Dis.* 1993; 168:195-201.



Medical

Pall Medical

25 Harbor Park Dr
Port Washington, NY 11050

516.484.3600 phone
medical.pall.com web
uscustomerservice@pall.com email

Europe

Pall Medical
a division of Pall Europe Ltd
Europa House, Havant Street
Portsmouth PO1 3PD, England

+44 (0)23 9230 3452 telephone
+44 (0)23 9230 3324 fax
Biosvc@Pall.com E-mail



Visit us on the Web at www.pall.com

International Offices

Pall Corporation has offices and plants throughout the world in locations such as: Argentina, Australia, Austria, Belgium, Brazil, Canada, China, France, Germany, Hong Kong, India, Indonesia, Ireland, Italy, Japan, Korea, Malaysia, Mexico, the Netherlands, New Zealand, Norway, Poland, Puerto Rico, Russia, Singapore, South Africa, Spain, Sweden, Switzerland, Taiwan, Thailand, the United Kingdom, the United States, and Venezuela. Distributors in all major industrial areas of the world.

The information provided in this literature was reviewed for accuracy at the time of publication. Product data may be subject to change without notice. For current information consult your local Pall distributor or contact Pall directly.

© 2006, Pall Corporation. Pall, , and Lipipor are trademarks of Pall Corporation. © indicates a registered trademark in the USA. **Filtration. Separation. Solution.**™ is a service mark of Pall Corporation.